

Quantifying the impact of early 21st century volcanic eruptions on global-mean surface temperature

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2017 Environ. Res. Lett. 12 054010

(<http://iopscience.iop.org/1748-9326/12/5/054010>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 138.63.210.70

This content was downloaded on 10/05/2017 at 14:30

Please note that [terms and conditions apply](#).

You may also be interested in:

[The role of external forcing and internal variability in regulating global mean surface temperatures on decadal timescales](#)

Lu Dong and Michael J McPhaden

[Observed and simulated temperature extremes during the recent warming hiatus](#)

Jana Sillmann, Markus G Donat, John C Fyfe et al.

[Internal variability of Earth's energy budget simulated by CMIP5 climate models](#)

M D Palmer and D J McNeall

[Effects of declining aerosols on projections of zonally averaged tropical precipitation](#)

L D Rotstayn, M A Collier and J-J Luo

[Examination of a climate stabilization pathway via zero-emissions using Earth system models](#)

Daisuke Nohara, J Tsutsui, S Watanabe et al.

[Quantifying the temperature-independent effect of stratospheric aerosol geoengineering on global-mean precipitation in a multi-model ensemble](#)

Angus J Ferraro and Hannah G Griffiths

[Rapid and extensive warming following cessation of solar radiation management](#)

Kelly E McCusker, Kyle C Armour, Cecilia M Bitz et al.

[Timing and seasonality of the United States 'warming hole'](#)

Nora R Mascioli, Michael Previdi, Arlene M Fiore et al.

[The global precipitation response to volcanic eruptions in the CMIP5 models](#)

Carley E Iles and Gabriele C Hegerl

Environmental Research Letters



LETTER

Quantifying the impact of early 21st century volcanic eruptions on global-mean surface temperature

OPEN ACCESS

RECEIVED
30 January 2017

REVISED
4 April 2017

ACCEPTED FOR PUBLICATION
12 April 2017

PUBLISHED
9 May 2017

Paul-Arthur Monerie¹, Marie-Pierre Moine, Laurent Terray and Sophie Valcke

CECI UMR 5318 CERFACS/CNRS, Toulouse, France

¹ Author to whom any correspondence should be addressed.

E-mail: pmonerie@gmail.com

Keywords: climate variability, hiatus, volcanic activity, climate modelling

Supplementary material for this article is available [online](#)

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Abstract

Despite a continuous increase in well-mixed greenhouse gases, the global-mean surface temperature has shown a quasi-stabilization since 1998. This muted warming has been linked to the combined effects of internal climate variability and external forcing. The latter includes the impact of recent increase in the volcanic activity and of solar irradiance changes. Here we used a high-resolution coupled ocean–atmosphere climate model to assess the impact of the recent volcanic eruptions on the Earth’s temperature, compared with the low volcanic activity of the early 2000s. Two sets of simulations are performed, one with realistic aerosol optical depth values, and the other with a fixed value of aerosol optical depth corresponding to a period of weak volcanic activity (1998–2002). We conclude that the observed recent increase in the volcanic activity led to a reduced warming trend (from 2003 to 2012) of 0.08 °C in ten years. The induced cooling is stronger during the last five-year period (2008–2012), with an annual global mean cooling of 0.04 °C (+/– 0.04 °C). The cooling is similar in summer (0.05 °C +/– 0.04 °C cooling) than in winter (0.03 °C +/– 0.04 °C cooling), but stronger in the Northern Hemisphere than in the Southern Hemisphere. Although equatorial and Arctic precipitation decreases in summer, the change in precipitation does not indicate robust changes at a local scale. Global heat content variations are found not to be impacted by the recent increase in volcanic activity.

1. Introduction and motivation

Despite a continuous increase in the concentration of well-mixed greenhouse gases (GHGs), the observed global mean air-surface temperature (GMST) trend has remained more or less steady from 2001 to 2013 (England *et al* 2014). This rate of temperature change does however not result in a significant slowdown of global warming rate with regard to the 1950–2013 time series (Rajaratnam *et al* 2015) and the pause was partly attributed to observational errors (Karl *et al* 2015). The recent observed GMST trend over the period of interest here (2003–2012) is at the lower end of the range of simulations performed in phase 5 of the Coupled Model Intercomparison Project Phase 5 (CMIP5; Taylor *et al* 2012) (Watanabe *et al* 2013). The CMIP5 climate models simulate a continuous increase in the rate of the temperature change (Kosaka and Xie 2013, Watanabe *et al* 2013, Meehl *et al* 2014). This has

raised the question of the role of decadal climate variability in reducing or exacerbating the impact of the global warming.

The surface temperature fluctuation can be mainly associated with the internal climate variability through a negative phase of the Pacific Decadal Oscillation (PDO) (Meehl *et al* 2011, 2014, Kosaka and Xie 2013, Watanabe *et al* 2013, 2014, Trenberth and Fasullo 2013, Guemas *et al* 2013, England *et al* 2014 and Douville *et al* 2015). During the last decade, the warming rate has been reduced, along with an increase in heat uptake, mainly in the Pacific and the Atlantic Ocean (Meehl *et al* 2011, Guemas *et al* 2013).

External forcing can also cause decadal-timescale changes in GMST. Some studies argue that the solar minimum around 2009 (Kopp and Lean 2011), the recent decrease in stratospheric water vapor concentration (Solomon *et al* 2010), the increase in emissions of anthropogenic sulfate aerosols (Kaufmann *et al* 2011,

Schmidt *et al* 2014) and the increase in stratospheric volcanic aerosol emissions (Solomon *et al* 2011, Fyfe *et al* 2013, Santer *et al* 2014, 2015, Haywood *et al* 2014, Ridley *et al* 2014, Schmidt *et al* 2014, Brühl *et al* 2015, Mills *et al* 2016) partly explain the reduction of the warming trend.

Systematic errors in certain external forcings in CMIP5 simulations of historical climate change explain the difference between simulated and observed warming rates during the ‘slowdown’ period. CMIP5 climate models are generally forced by unrealistically low stratospheric aerosols optical depth after 2000 (Fyfe *et al* 2013, among others). The cooling effect of the solar activity was also neglected (Schmidt *et al* 2014). Biases in simulating El Niño Southern Oscillation phases also lead to GMST overestimation (Schmidt *et al* 2014). Observational errors (Karl *et al* 2015) and model-observation comparison biases (Cowtan *et al* 2015) can also be partly responsible for the apparent gap between models and observations. Richardson *et al* (2016) have therefore shown that the gap between model simulations and observations are greatly reduced when model outputs are processed the same way as the HadCRUT4 observations over the oceans, and in taking into account the lack of values over the Arctic in the observations. HadCRUT4 is a dataset of observed near-surface air temperature (Morice *et al* 2012).

Here we study the recent surface temperature fluctuation by analyzing the impact of the volcanic eruptions that have occurred between 2003 and 2012. In order to simulate the impact of the recent volcanic activity on climate we performed two sets of sensitivity experiments, with two different data sets of stratospheric aerosol optical depth (SAOD).

Our approach is similar to that followed by Solomon *et al* (2011), Fyfe *et al* (2013), Ridley *et al* (2014) and Haywood *et al* (2014), who analyzed the impact of the recent volcanic eruptions on the GMST. These authors have concluded that the recent volcanic eruptions may have cooled GMST by 0.02 °C (Haywood *et al* 2014) to 0.05 °C–0.12 °C (Ridley *et al* 2014). Solomon *et al* (2011) and Fyfe *et al* (2013) found an intermediate values of 0.05 °C–0.07 °C.

The main objective of Fyfe *et al* 2013 was to explore the consequences of systematic errors in model representation of early 21st century volcanic aerosol forcing (using a near-zero radiative forcing as a baseline). Here we prefer using an alternative approach with an SAOD baseline set at a fixed value, deduced from a weak volcanic activity period (1998–2002). We address the impact of the recent increase in the volcanic activity, in comparison to a period of a low-volcanic activity as a baseline. We mainly assess the impact of the moderate strengthening in the volcanic activity on a recent decade (2003–2012) rather than documenting the impact of using biased low forcing. To analyze the impact of natural volcanic activity variability in (the absence of

strong volcanic events) we have to deal with weaker differences in SAOD, than in previous studies (Fyfe *et al* 2013, among others). The impact of such a low change in radiative forcing is thus still an open question. The question of the misrepresentation of the volcanic external forcing in global climate models is therefore indirectly addressed.

Our scientific question: What is the impact, on the GMST, of the recent of moderate volcanic activity (2003–2012) compared to a period of low volcanic activity, such as from 1998–2002?


2. Data, model and simulations

2.1. The CERFACS-HR climate model and simulations

We used the CERFACS-HR climate model, a high resolution coupled climate model developed at CERFACS (Centre Européen de Recherche et de Formation Avancée en Calcul Scientifique). The ocean component is the Nucleus for European Models of the Ocean (NEMO) v3.4 (Madec 2008) model on an ORCA025L75 grid (horizontal resolution of 0.25° and 75 vertical levels). The atmospheric model is ARPEGE-climate v5.3 developed at Meteo-France/CNRM (Centre National de Recherches Météorologiques) with a horizontal resolution of about 50 kilometers and 31 vertical levels (T359L31). More information on CERFACS-HR is given in Monerie *et al* (2017, hereafter noted MO17).

The coupled model is initialized with the GLORYS2V1 ocean reanalysis (Ferry *et al* 2012) and the atmosphere initial state is given by a sea-surface temperature (SST)-forced ARPEGE simulation. The simulations are full-field initialized in 2002 (in November). We performed six realizations, which differ by their atmospheric initial state, corresponding to different days in November. All simulations are forced by the estimated historical values in GHGs, solar irradiance, black carbon, particulate organic matter, dust, sea salt and sulfate aerosols. The optical depths of tropospheric aerosols are given by Szopa *et al* (2013).

The ability of CERFACS-HR to simulate climate is provided in MO17. Before the drift correction, simulations exhibit large positive biases of SST over the Austral Ocean and over the equatorial Pacific and Atlantic Ocean (up to 4 °C), the Arctic experiences a cold bias of 5 °C. These biases are commonly obtained with climate model simulations. After drift correction CERFACS-HR is able to simulate and to predict the global mean surface temperature for the upcoming five years and exhibits high skill in predicting SST over the North Atlantic, equatorial Atlantic and the Indian Oceans (MO17). CERFACS-HR has also high skill in predicting decadal trends in Arctic sea-ice extent and volume. Moreover CERFACS-HR has been used in a previous study to perform a suite of decadal hindcasts



(10 year hindcasts starting each year from November 1993 to November 2012) that reproduce a reduced surface warming rate during the recent decades (MO17). In these hindcast simulations, the 2003–2012 trend in GMST is associated with a negative phase of the PDO, and is therefore mainly explained by internal climate variability. This result is likely to be linked to the model initial state, since the PDO and the Interdecadal Pacific Oscillation (IPO) shifted from a positive to a negative phase in the early 2000s (Trenberth and Fasullo 2013, England *et al* 2014). This issue is discussed later, in section 3.4. However, MO17 carried an important limitation, i.e. an unrealistic evolution of the volcanic eruptions due to the use of a background condition in SAOD instead of realistic SAOD data. Here we analyze the added

value of a realistic representation of the volcanic activity.

To meet this objective, two sets of simulations are performed:

- The first set, hereafter referred as VER, includes six members using the SAOD data of Vernier *et al* (2011), therefore taking into account the recent evolution of the volcanic activity (figure 1 (a), red line).
- The second set, hereafter referred as BAK, is realized by performing also an ensemble of six members using a background value of SAOD, based on the 1998–2002 lower values of Vernier *et al* (2011) (figure 1(a), blue line). This baseline

allows quantifying the impact of the recent volcanic activity in comparison with a period of minimal volcanic activity. Here we do not document the consequences on the climate simulations of using a biased low volcanic forcing (with a near-zero value in SAOD), as done in many studies (Solomon *et al* 2011, Fyfe *et al* 2013, Ridley *et al* 2014). The background in SAOD is however never equal to zero (as highlighted in Solomon *et al* 2011) and our choice of the baseline value of SAOD is more adapted to our objective.

The impact of the recent volcanic eruptions is quantified by computing the VER-BAK differences.

Climate models do not perfectly simulate the observed climate. When initialized with observations models drift toward their preferred imperfect climatology (Meehl *et al* 2009). This drift leads to biases in hindcasts and has to be removed. We used an additional set of ten 10 year hindcasts, initialized every year from 1993 to 2002 (5 members \times 10 start dates), described in MO17, to estimate a drift correction, following the World Climate Research Program recommendations (ICPO 2011). The drift was then removed from VER and BAK at each lead-time (time step after the model initialization) (see supplementary materials available at stacks.iop.org/ERL/12/054010/mmedia for further details on the bias adjustment method). Assuming that the drift is equal in the two sets of simulations, we did not remove the drift prior to computing the VER-BAK differences (in what follows, we only remove the drift when showing separately VER and BAK results). The VER-BAK differences are thus computed at the CERFACS-HR horizontal resolution. We checked that the result is not dependent on the removal of the drift prior to compute the anomalies (not shown).

2.2. Data

We compare the simulated GMST to several data sets of near-surface air temperature:

- i. ERA-Interim reanalysis (hereafter noted ERAI; Dee *et al* 2011), available from 1979 to present on a $1.5^\circ \times 1.5^\circ$ horizontal resolution;
- ii. the Cowtan and Way (2014) data, which consists in an improvement of HadCRUT4 (Morice *et al* 2012) through a better infilling of poorly sampled regions; this data set, provided at a $5^\circ \times 5^\circ$ horizontal resolution is available from 1850 to present;
- iii. The NASA's Goddard Institute for Space Studies Surface Temperature Analysis (GISTEMP) smoothed at 250 km (Hansen *et al* 2010), available from 1880 to present at a $2^\circ \times 2^\circ$ horizontal resolution;
- iv. The NOAA Merged Land–Ocean Surface Temperature (MLOST) version 3b of the National Climatic Data Center (NCDC), available from 1880 to present at a $5^\circ \times 5^\circ$ horizontal resolution (Vose *et al* 2012);
- v. The Berkeley Earth Surface Temperature dataset (BEST), which is an interpolated dataset of surface air temperature anomalies (with respect to the 1951–1980 period) over land and ocean built from temperature records by using statistical techniques (Rohde *et al* 2013); BEST is available from 1850 to present at a $1^\circ \times 1^\circ$ horizontal resolution.

3. Results

3.1. Radiative forcing

The 10 year trends of the top of the atmosphere (TOA) net shortwave, longwave and net radiative fluxes are shown in figures 1(b)–(d). The inter-member spread is large in both BAK and VER simulations. We found a large interannual variability in TOA for both BAK, VER, we assume that this can lead to large inter-member spread over short-periods (ten years). As the six-simulation ensemble average is strongly influenced by outliers, we also show the ensemble median. Both BAK and VER exhibit negative 10 year trends in TOA net shortwave (figure 1(b)). The VER trend is stronger than the BAK trend, indicating a stronger decrease in the net shortwave radiation. Moreover the VER-BAK difference is greater than 1 BAK standard deviation (the standard deviation obtained from the six realizations of BAK, figure 1(b)). We split the net shortwave flux into its incoming and outgoing parts (not shown). As the former is the same in both simulations, this indicates that recent eruptions increased the outgoing shortwave radiation, due to the volcanic cloud albedo effect. Santer *et al* (2014) noted a similar observed change in TOA net shortwave, with a decreased radiative forcing of 0.25 W m^{-2} per decade over January 2001–December 2012 (the median of the VER-BAK difference is of 0.25 W m^{-2} per decade between 2003 and 2012).

The TOA longwave flux trends are positive (i.e. downward) and stronger in VER than in BAK (figure 1(c)). This denotes a stronger absorption of upward longwave fluxes in VER, which lead to a warming of the troposphere. It is not strong since the differences in the mean and the median are weaker than 1 BAK standard deviation. The VER minus BAK difference in TOA longwave is due to volcanic eruptions that inject sulphur dioxide (and to a lesser extent, hydrogen sulfide) into the stratosphere, which are converted into sulfate aerosols that act to both scatter solar radiation and absorb longwave and near-infra-red radiation, as shown in Robock (2000) and Timmreck (2012).

Finally the VER simulations exhibit a more negative (i.e. upward) trend in net TOA heat fluxes

Table 1. 10 year GMST trend for the different observations, ERAI and the modeled 10 year GMST trend. The 2003–2012 GMST trend error is given by half the 2.5%–97.5% regression coefficient confidence interval. For VER and BAK, the inter-member spread in GMST trend is added.

Data	2003–2012 GMST trend	2003–2012 GMST trend error
ERAI	−0.005 °C/decade	+/− 0.205 °C/decade
Cowtan and Way	+0.028 °C/decade	+/− 0.166 °C/decade
MLOST	−0.044 °C/decade	+/− 0.115 °C/decade
GISTEMP	−0.038 °C/decade	+/− 0.130 °C/decade
BEST	+0.016 °C/decade	+/− 0.172 °C/decade
VER	+0.037 °C/decade +/− 0.074 °C/decade	+/− 0.064 °C/decade
BAK	+0.118 °C/decade +/− 0.053 °C/decade	+/− 0.077 °C/decade

than BAK and the difference between VER-BAK is even more robust when considering the medians (figure 1(d); online supplementary table S1). This difference indicates a stronger increase in outgoing heat fluxes in VER compared to BAK. The difference, of 0.21 W m^{-2} in ten years (median) is consistent with the modelled studies of Brühl *et al* (2015) who obtained a decrease in the radiative forcing by up to 0.2 W m^{-2} due to the volcanic eruptions between 2005 and 2011.

It is therefore expected that VER simulates a colder mean climate than BAK, which is consistent with the known impact of volcanic eruptions on radiative fluxes at the top of Earth's atmosphere.

3.2. Global mean surface temperature

The 2003–2012 GMST trend ranges from a weak warming in BEST and in Cowtan and Way to a weak cooling in ERAI, in MLOST and in GISTEMP (table 1). Observed 10 year trends have large uncertainties for all data sets (third column of the table 1). The observed trends are close to the 1998–2012 trend ($0.03 \text{ °C per decade}$), a result found by Kosaka and Xie (2015) with the HadCRUT 4 dataset.

The observed data sets have different horizontal resolutions and use various infilling and coverage over the polar regions. Hansen *et al* (2010) and Cowtan and Way (2014) have shown that discrepancies across the datasets in GMST are mainly due to the estimated temperature in polar regions, where observations are sparse.

The BAK 2003–2012 trend is warmer than the corresponding ERAI trend, with a warming of $+0.12 \text{ °C (+/− 0.05 °C)}$ in ten years (close to the observed GMST 1951–2012 trend, of $+0.11 \text{ °C per decade}$, according to Kosaka and Xie 2015) (table 1 and figure 2(a)). In VER, the 2003–2012 trend is also positive ($+0.04 \text{ °C (+/− 0.07 °C per decade)}$). This trend is close to BEST and 'Cowtan and Way', which are the most reliable observations for the analysis of the GMST, due to a better representation of the surface temperature at high latitudes (Dodd *et al* 2015). The difference between VER and BAK trend is determined from the time series of paired differences between VER and BAK. It reduces noise levels by subtracting variability components common to VER and BAK. The BAK-VER trend is statistically significant at the

95% confidence level according to a Spearman's rank correlation test. This suggests that the recent volcanism activity has a significant impact on the GMST, reducing the warming trend by about 0.08 °C in ten years, compared to a period of weak volcanic activity.

BAK and VER GMST differences get stronger with time (figure 2(a)). This can be due to the increase in SAOD after 2007 with stronger peaks mainly due to the Tavurvur (in 2007), Sarychev (in 2009) and Nabro (in 2011) eruptions. From 2003 to 2005, the SAOD values in VER are close to the background (figure 1).

In figure 2 we chose ERAI as our observation data set since it provides information for a large set of variables (see the additional analyses in the supplementary materials) with a complete Earth coverage (avoiding model-observation differences due to different spatial coverages, such as missing values over the Arctic as reported in Hansen *et al* 2010 and Cowtan and Way 2014). Note that ERAI can be considered as a reference, since estimates of Arctic temperatures and temperature trends are realistic (Simmons *et al* 2014, Simmons and Poli 2015). In figures 2 (b)–(d), we show the VER-BAK GMST differences for the 5 last years of the simulations (2008–2012), when the difference between the VER and BAK AOD files is the strongest and when the observed impact of the volcanic activity is the strongest since 2003 (Santer *et al* 2015). The global impact is of $-0.04 \text{ °C (+/− 0.04 °C)}$. The difference is similar over land ($-0.05 \text{ °C (+/− 0.06 °C)}$) than over the oceans ($-0.04 \text{ °C (+/− 0.03 °C)}$) (figures 2(c),(d)). All these differences are statistically significant at a 95% confidence level following a Monte Carlo approach (more details are available in the supplementary materials; table S1 and table S2). VER simulates a colder mean climate than BAK for the global Earth (figure 2(b)), the land (figure 2(c)) and the ocean (figure 2(d)).

The ERAI-VER difference is however stronger than the VER-BAK difference (and stronger than 1 VER standard deviation on global average). There are clearly other phenomena to take into account to reproduce the observed slightly negative trend, such as, for instance, the cooling over Eurasia in winter, which arises essentially from atmospheric internal variability (Li *et al* 2015).




Figure 2. (a) Global-mean surface temperature evolution for ERAI (gray line), BAK (blue line) and VER (red line); the corresponding 2003–2012 linear trend is displayed with discontinuous lines. The blue (red) shading represents the spread (\pm one standard deviation) of the BAK (VER) ensemble-members. Five year (2008–2012) mean in (b) global-mean surface temperature, (c) mean temperature over land and (d) ocean for VER, BAK and ERAI. Circles represent each BAK and VER simulations, crosses represent the ensemble-mean and triangles the ensemble-median. The gray shading (discontinuous lines) is used to delimit the spread (\pm one standard deviation around the ensemble mean) of the BAK (VER) ensemble.


3.3. Regional response in temperature, precipitation and heat content

Haywood *et al* (2014) have shown that the impact of the recent volcanic eruptions is not spatially homogeneous, for example stronger in the Northern Hemisphere than in the Southern Hemisphere. In this section, we focus on the regional responses in temperature and precipitation. Moreover we explore the seasonality of the response.

From November to April (NDJFMA, which includes the boreal winter) the difference in GMST is of $-0.03\text{ }^{\circ}\text{C}$ ($\pm 0.04\text{ }^{\circ}\text{C}$). The impact is significant only over several restricted areas (figure 3(a)). Precipitation change only exhibits sparse significant anomalies (figure 3(c)).

Impacts are of comparable magnitude from May to October (MJJASO, which encompasses the boreal summer) with a decrease in GMST of $0.05\text{ }^{\circ}\text{C}$

($\pm 0.04\text{ }^{\circ}\text{C}$). The zonally-averaged temperature decreases over the northern hemisphere, due to a cooling of the Arctic. The subpolar gyre warms and the tropical Atlantic and eastern Pacific ocean cool down (figure 3(b)). A stronger impact over the Arctic is consistent with Fyfe *et al* (2013) and Haywood *et al* (2014). Moreover the cooling over the Arctic is stronger in MJJASO than in NDJFMA, since it receives more solar flux in summer than in winter, allowing a more efficient direct effect of the volcanic forcing. Precipitation decreases over the equator and south of it, as also suggested in Fyfe *et al* (2013). This result however disagrees with Haywood *et al* (2014), who have found a southward shift of the tropical precipitation due to a cooling of the subtropical Atlantic Ocean. The precipitation pattern consists in an increase in precipitation over the western Pacific Ocean and a decrease over the western coast of South



America, along the equator, which is consistent with a La Niña pattern.

It has been shown that major volcanic eruptions can affect the ENSO variability, leading to an El Niño (Ohba *et al* 2013) or a La Niña (Maher *et al* 2015) event several months after the eruption. Here there is no robust change in ENSO variability (not shown).

The volcanic eruptions can also impact the North Atlantic Oscillation (NAO) variability (Shindell *et al* 2004). We obtain a tripole in sea-level pressure (SLP) anomaly: a positive pressure anomaly north of Europe and negative ones west of Spain and over north-eastern Canada, i.e. that project on a negative phase of NAO (online supplementary figure S1). The negative NAO pattern is consistent with a decrease in surface zonal wind speed over the North Atlantic (figure S1), and the warming of the north Atlantic SST. However, a NAO index, derived from our simulations, does not indicate a robust change in the NAO variability (not shown).

Gleckler *et al* (2016) argued that the recent volcanic eruptions led to a decrease in ocean heat-uptake. We however found no strong differences between BAK and VER when considering the global mean heat-content surface to 700 m, 700 m to 2000 m and the deep ocean (>2000 m) (online supplementary figure S2). Here the heat uptake increase is weak and only located to the eastern equatorial Pacific Ocean (online supplementary figure S3).

3.4. Impact of the initial condition and mean state

Pohlmann *et al* (2004) found that the North Atlantic SSTs, Nordic Seas and Southern Ocean exhibit predictability on multidecadal time scales, owing to

the model initialization and the oceanic inertia. Numerous predictability studies have highlighted the importance of the initialization to predict the GMST (Bellucci *et al* 2013, García-Serrano *et al* 2015, Karspeck *et al* 2015). Here we analyze the 10 year trend in GMST for both BAK and VER. In both simulations, temperature increases over the subtropical Pacific Ocean and decreases over the eastern Pacific Ocean, exhibiting a negative phase of the PDO (online supplementary figure S4). CERFACS-HR systematically reproduces a negative phase of the PDO over the 2003–2012 period, as also shown in MO17 due to the ocean initialization (these simulations have common ocean initial conditions).

On the one hand the predictability of the SSTs is mainly given by the model initialization. On the other hand the mean state of the climate system may modulate the impact of the volcanic eruptions (Zanchettin *et al* 2013). We can thus hypothesize that the impact of the volcanic eruptions could be different without the simulated negative phase of the PDO, and the associated abnormally low GMST.

4. Discussion/conclusion

During the years 2003 to 2012, moderate volcanic eruptions led to an increase in stratospheric AOD (Vernier *et al* 2011, Santer *et al* 2013) that impacted the GMST. This phenomenon explains a part of the recent temperature fluctuation. We used the Vernier *et al* (2011) AOD, which takes into account the presence of the recent volcanic eruptions. Our results suggest, in agreement with Solomon *et al* (2011), Fyfe *et al*

(2013), Haywood *et al* (2014), Schmidt *et al* (2014), Ridley *et al* (2014), Santer *et al* (2014, 2015), Brühl *et al* (2015) and Mills *et al* (2016) that the recent volcanic activity has reduced the GMST and is therefore one of the causes of the recent reduced warming trend.

A cooling of $0.04\text{ }^{\circ}\text{C}$ ($\pm 0.04\text{ }^{\circ}\text{C}$) over the 2008–2012 period was found in this study. Such result is consistent with Haywood *et al* (2014), in which the decrease of temperature is of $0.02\text{ }^{\circ}\text{C}$ – $0.03\text{ }^{\circ}\text{C}$. A stronger response found in the Northern Hemisphere than the Southern Hemisphere is also consistent with this author.

Both BAK and VER simulations exhibit a negative phase of the PDO, which is therefore not the only mechanism responsible for a global warming rate slowdown. For instance, even limited increase in volcanism activity, as it occurred in the last decade, has also exacerbated the eastern Pacific cooling.

The induced cooling ($0.04\text{ }^{\circ}\text{C}$) is weaker than in Solomon *et al* (2011) and Fyfe *et al* (2013), who obtained a cooling of $0.07\text{ }^{\circ}\text{C}$. In addition to the sensitivity to volcanic eruptions, several factors can explain this difference. (i) We did not use the same SAOD baseline as in Haywood *et al* (2014), Solomon *et al* (2011) and Fyfe *et al* (2013). (ii) The result can be model-dependent. (iii) Ocean initial conditions can play an important role in the model response and (iv) the relatively small sample size may also lead to large uncertainties.

Fyfe *et al* (2013), Haywood *et al* (2014), Santer *et al* (2014), Ridley *et al* (2014) and Schmidt *et al* (2014) used a near-zero value in volcanic forcing to document the importance of volcanic forcing biases. Here we used a baseline based on a weak volcanic activity (a more realistic value since the volcanic activity is not expected to stop) and are dealing with weaker anomalies in radiative forcing. The obtained cooling is thus weaker than in the aforementioned studies. We however assume that this experimental protocol is noteworthy since we can document the impact of natural volcanic activity variability, in absence of intense volcanic events, rather than documenting the impact of using biased low forcing.

Vernier *et al* (2011) SAOD neglects substantial amounts of volcanic aerosols below 15 km and therefore underestimates total radiative forcing due to the recent eruptions (Ridley *et al* 2014 and Mills *et al* 2016). The impact of including the recent volcanic eruptions in climate models could thus have been stronger than previously highlighted and obtained here. Ridley *et al* (2014) estimated a global cooling of 0.05 to $0.12\text{ }^{\circ}\text{C}$ since 2000 with a better estimation of the global volcanic aerosol forcing.

Results obtained in this study show that moderate volcanic eruptions cause a small cooling of the Earth and highlight the importance of taking into account a realistic volcanic forcing in the climate models. However, this volcano-related cooling is not sufficient

to fully explain the recent surface temperature stagnation. The cooling generated by volcanic eruptions is thus not the unique cause for the recent decrease in the overall trend of temperature increase.

The impact of the recent volcanic eruptions is a promising topic for understanding the GMST evolution and its better understanding can help to improve climate model simulations since the stratospheric AOD are not well represented by the CMIP5 climate models in the recent decades (Fyfe *et al* 2013). Such a study should however be extended to a multi-model analysis, in order to assess the robustness of the results with regard to model sensitivity, and/or with different ocean initial conditions.

Acknowledgments

We thank the three anonymous reviewers for their helpful and constructive suggestions and comments. The authors gratefully acknowledge the support from the Seventh Framework Programme (FP7) of the European Commission (grant agreement 308378) SPECS Project, the PRECLIDE project funded by the BNP-PARIBAS foundation. Simulations were run thanks to PRACE HiResClim I and II projects.

References

- Bellucci A, Gualdi S, Masina S, Storto A, Scoccimarro E, Cagnazzo C, Fogli P, Manzini E and Navarra A 2013 Decadal climate predictions with a coupled OAGCM initialized with oceanic reanalyses *Clim. Dynam.* **40** 1483–97
- Brühl C, Lelieveld J, Tost H, Höpfner M and Glatthor N 2015 Stratospheric sulfur and its implications for radiative forcing simulated by the chemistry climate model EMAC *J. Geophys. Res. Atmos.* **120** 2103–18
- Cowan K and Way R G 2014 Coverage bias in the HadCRUT4 temperature series and its impact on recent temperature trends *Q. J. R. Meteorolog. Soc.* **140** 1935–44
- Cowan K, Hausfather Z, Hawkins E, Jacobs P, Mann M E, Miller S K, Steinman B A, Stolpe M B and Way R G 2015 Robust comparison of climate models with observations using blended land air and ocean sea surface temperatures *Geophys. Res. Lett.* **42** 6526–34
- Dee D P *et al* 2011 The ERA-Interim reanalysis: configuration and performance of the data assimilation system *Q. J. R. Meteorolog. Soc.* **137** 553–97
- Dodd E M, Merchant C J, Rayner N A and Morice C P 2015 An investigation into the impact of using various techniques to estimate Arctic surface air temperature anomalies *J. Clim.* **28** 1743–63
- Douville H, Volodrova A and Geoffroy O 2015 The recent global warming hiatus: what is the role of Pacific variability? *Geophys. Res. Lett.* **42** 880–8
- England M H *et al* 2014 Recent intensification of wind-driven circulation in the Pacific and the ongoing warming hiatus *Nat. Clim. Change* **4** 222–7
- Ferry N *et al* 2012 GLORYS2V1 global ocean reanalysis of the altimetric era (1993–2009) at meso scale *Mercator Ocean–Q. Newslett.* **44**
- Fyfe J C, von Salzen K, Cole J N S, Gillett N P and Vernier J P 2013 Surface response to stratospheric aerosol changes in a coupled atmosphere–ocean model *Geophys. Res. Lett.* **40** 584–8

- García-Serrano J, Guemas V and Doblas-Reyes F J 2015 Added-value from initialization in predictions of Atlantic multi-decadal variability *Clim. Dynam.* **44** 2539–55
- Gleckler P J, Durack P J, Stouffer R J, Johnson G C and Forest C E 2016 Industrial-era global ocean heat uptake doubles in recent decades *Nat. Clim. Change* **6** 394–8
- Guemas V, Doblas-Reyes F J, Andreu-Burillo I and Asif M 2013 Retrospective prediction of the global warming slowdown in the past decade *Nat. Clim. Change* **3** 649–53
- Hansen J, Ruedy R, Sato M and Lo K 2010 Global surface temperature change *Rev. Geophys.* **48** RG4004
- Haywood J M, Jones A and Jones G S 2014 The impact of volcanic eruptions in the period 2000–2013 on global mean temperature trends evaluated in the HadGEM2-ES climate model *Atmos. Sci. Lett.* **15** 92–6
- ICPO 2011 Data and bias correction for decadal climate predictions *International CLIVAR Project Office Publication, Series 150* (www.wcrp-climate.org/decadal/references/DCPP_Bias_Correction.pdf) 5 pp
- Kaufmann R K, Kauppi H, Mann M L and Stock J H 2011 Reconciling anthropogenic climate change with observed temperature 1998–2008 *Proc. Natl Acad. Sci. U.S.A.* **108** 11790–3
- Karl T R, Arguez A, Huang B, Lawrimore J H, McMahon J R, Menne M J, Peterson T C, Vose R S and Zhang H-M 2015 Possible artifacts of data biases in the recent global surface warming hiatus *Science* **348** 1469–72
- Karspeck A, Yeager S, Danabasoglu G and Teng H 2015 An evaluation of experimental decadal predictions using CCSM4 *Clim. Dynam.* **44** 907–23
- Kopp G and Lean J L 2011 A new, lower value of total solar irradiance: evidence and climate significance *Geophys. Res. Lett.* **38** L01706
- Kosaka Y and Xie S-P 2013 Recent global-warming hiatus tied to equatorial Pacific surface cooling *Nature* **501** 403–7
- Kosaka Y and Xie S-P 2015 Tropical Pacific influence on the recent hiatus in surface global warming US CLIVAR, 10
- Li C, Stevens B and Marotzke J 2015 Eurasian winter cooling in the warming hiatus of 1998–2012 *Geophys. Res. Lett.* **42** 8131–9
- Madec G 2008 NEMO ocean engine *Note du Pole de modélisation* 27 (France: Institut Pierre-Simon Laplace (IPSL))
- Maher N, McGregor S, England M H and Gupta A S 2015 Effects of volcanism on tropical variability *Geophys. Res. Lett.* **42** 6024–33
- Meehl G A *et al* 2009 Decadal prediction *Bull. Am. Meteorol. Soc.* **90** 1467–85
- Meehl G A, Teng H and Arblaster J M 2014 Climate model simulations of the observed early-2000s hiatus of global warming *Nat. Clim. Change* **4** 898–902
- Meehl G A, Arblaster J M, Fasullo J T, Hu A and Trenberth K E 2011 Model-based evidence of deep-ocean heat uptake during surface-temperature hiatus periods *Nat. Clim. Change* **1** 360–4
- Mills M J *et al* 2016 Global volcanic aerosol properties derived from emissions, 1990–2014, using CESM1(WACCM) *J. Geophys. Res. Atmos.* **121** 2332–48
- Monerie P A, Coquart L, Maisonnave É, Moine M P, Terray L and Valcke S 2017 Decadal prediction skill using a high-resolution climate model *Clim. Dynam.* (<https://doi.org/10.1007/s00382-017-3528-x>)
- Morice C P, Kennedy J J, Rayner N A and Jones P D 2012 Quantifying uncertainties in global and regional temperature change using an ensemble of observational estimates: the HadCRUT4 data set *J. Geophys. Res. Atmos.* **117** D08101
- Ohba M, Shioyama H, Yokohata T and Watanabe M 2013 Impact of strong tropical volcanic eruptions on ENSO simulated in a coupled GCM *J. Clim.* **26** 5169–82
- Pohlmann H, Botzet M, Latif M, Roesch A, Wild M and Tschuck P 2004 Estimating the decadal predictability of a coupled AOGCM *J. Clim.* **17** 4463–72
- Rajaratnam B, Romano J, Tsiang M and Diffenbaugh N S 2015 Debunking the climate hiatus *Clim. Change* **133** 129–40
- Richardson M, Cowtan K, Hawkins E and Stolpe M B 2016 Reconciled climate response estimates from climate models and the energy budget of Earth *Nat. Clim. Change* **6** 931–5
- Ridley D A *et al* 2014 Total volcanic stratospheric aerosol optical depths and implications for global climate change *Geophys. Res. Lett.* **41** 7763–9
- Robock A 2000 Volcanic eruptions and climate *Rev. Geophys.* **38** 191–219
- Rohde R, Muller R, Jacobsen R, Perlmutter S, Rosenfeld A, Wurtele J, Curry J, Wickham C and Mosher S 2013 Berkeley earth temperature averaging process *Geoinfor. Geostat. An Overview* 1 1–13
- Santer B D *et al* 2014 Volcanic contribution to decadal changes in tropospheric temperature *Nat. Geosci.* **7** 185–9
- Santer B D *et al* 2015 Observed multivariable signals of late 20th and early 21st century volcanic activity *Geophys. Res. Lett.* **42** 500–9
- Schmidt G A, Shindell D T and Tsigaridis K 2014 Reconciling warming trends *Nat. Geosci.* **7** 158–60
- Shindell D T, Schmidt G A, Mann M E and Faluvegi G 2004 Dynamic winter climate response to large tropical volcanic eruptions since 1600 *J. Geophys. Res. Atmos.* **109** D05104
- Simmons A J and Poli P 2015 Arctic warming in ERA-Interim and other analyses *Q. J. R. Meteorol. Soc.* **141** 1147–62
- Simmons A J, Poli P, Dee D P, Berrisford P, Hersbach H, Kobayashi S and Peubey C 2014 Estimating low-frequency variability and trends in atmospheric temperature using ERA-Interim *Q. J. R. Meteorol. Soc.* **140** 329–53
- Solomon S, Daniel J S, Neely R R, Vernier J-P, Dutton E G and Thomason L W 2011 The persistently variable ‘background’ stratospheric aerosol layer and global climate change *Science* **333** 866–70
- Solomon S, Rosenlof K H, Portmann R W, Daniel J S, Davis S M, Sanford T J and Plattner G-K 2010 Contributions of stratospheric water vapor to decadal changes in the rate of global warming *Science* **327** 1219–23
- Szopa S *et al* 2013 Aerosol and ozone changes as forcing for climate evolution between 1850 and 2100 *Clim. Dynam.* **40** 2223–50
- Taylor K E, Stouffer R J and Meehl G A 2012 An overview of CMIP5 and the experiment design *Bull. Am. Meteorol. Soc.* **93** 485–98
- Timmreck C 2012 Modeling the climatic effects of large explosive volcanic eruptions *Wiley Interdiscip. Rev. Clim. Change* **3** 545–64
- Trenberth K E and Fasullo J T 2013 An apparent hiatus in global warming *Earth’s Future* **1** 19–32
- Vernier J P *et al* 2011 Major influence of tropical volcanic eruptions on the stratospheric aerosol layer during the last decade *Geophys. Res. Lett.* **38** L12807
- Vose R S *et al* 2012 NOAA’s merged land–ocean surface temperature analysis *Bull. Am. Meteorol. Soc.* **93** 1677–85
- Watanabe M, Shioyama H, Tatebe H, Hayashi M, Ishii M and Kimoto M 2014 Contribution of natural decadal variability to global warming acceleration and hiatus *Nat. Clim. Change* **4** 893–7
- Watanabe M, Kamae Y, Yoshimori M, Oka A, Sato M, Ishii M, Mochizuki T and Kimoto M 2013 Strengthening of ocean heat uptake efficiency associated with the recent climate hiatus *Geophys. Res. Lett.* **40** 3175–9
- Zanchettin D, Bothe O, Graf H F, Lorenz S J, Luterbacher J, Timmreck C and Jungclaus J H 2013 Background conditions influence the decadal climate response to strong volcanic eruptions *J. Geophys. Res. Atmos.* **118** 4090–106